

# Big Bounce and Inflation from Spin and Torsion

Nikodem Popławski



Institute for Theory and Computation Luncheon  
Harvard-Smithsonian Center for Astrophysics  
September 29, 2016

**Cosmic Microwave Background**

**Dark Energy  
Accelerated Expansion**

**Afterglow Light  
Pattern  
380,000 yrs.**

**Dark Ages**

**Development of  
Galaxies, Planets, etc.**

**Inflation**


**Big Bang**

**Quantum  
Fluctuations**

**1st Stars  
about 400 million yrs.**

**Big Bang Expansion**

**13.7 billion years**



# Problems of general relativity

General relativity describes gravity as curvature of spacetime.

Singularities: points with infinite density of matter.

Incompatible with quantum mechanics. We need quantum gravity. It may resolve the singularity problem.

Field equations contain the conservation of orbital angular momentum, contradicting Dirac equation which gives the conservation of total angular momentum (orbital + spin) and allows spin-orbit exchange.

Simplest extension of GR to include QM spin: **Einstein-Cartan theory**. It also resolves the singularity problem.

# Problems of big-bang cosmology & inflation

Big-bang singularity.

What caused the big bang? What existed before?

Inflation (exponential expansion of the early Universe) solves the flatness and horizon problems, and predicts the observed spectrum of CMB perturbations. What caused inflation?  
(scalar fields are usually used)

Why did inflation end? (no eternal inflation)

Einstein-Cartan theory replaces the singular Big Bang by a non-singular **Big Bounce**. The dynamics immediately after the bounce explains the flatness/horizon problems. [NP, PLB 694, 181 \(2010\)](#).

# Einstein-Cartan-Sciama-Kibble gravity

Spacetime has curvature and **torsion**.

$$S^k{}_{ij} = \Gamma_{[ij]}^k$$

Cartan equations:

**Torsion** is proportional to the **spin** density of fermions. ECSK in vacuum reduces to GR and passes all observational tests. It differs from GR at densities around  $10^{45}$  kg/m<sup>3</sup> and higher.

$$S_{jik} - S_i g_{jk} + S_k g_{ji} = -\frac{1}{2} \kappa S_{ikj}$$

[arXiv.org > gr-qc > arXiv:0911.0334](https://arxiv.org/abs/gr-qc/0911.0334)

Einstein equations:

**Curvature** is proportional to the **energy and momentum** density.

$$G^{ik} = \kappa T^{ik} + \frac{1}{2} \kappa^2 \left( s^{ij}{}_j s^{kl}{}_l - s^{ij}{}_l s^{kl}{}_j - s^{ijl} s^k{}_{jl} + \frac{1}{2} s^{jli} s_{jl}{}^k + \frac{1}{4} g^{ik} (2s_j{}^l{}_m s^{jm}{}_l - 2s_j{}^l{}_l s^{jm}{}_m + s^{jlm} s_{jlm}) \right)$$

# Universe with spin fluid

Dirac particles can be averaged macroscopically as a spin fluid.

$$s^{\mu\nu\rho} = s^{\mu\nu}u^\rho \quad s^{\mu\nu}u_\nu = 0 \quad s^2 = s^{\mu\nu}s_{\mu\nu}/2$$

Einstein-Cartan equations for a (closed) FLRW Universe become Friedman equations.

$$\begin{aligned} \dot{a}^2 + 1 &= \frac{1}{3}\kappa\left(\epsilon - \frac{1}{4}\kappa s^2\right)a^2, \\ \dot{a}^2 + 2a\ddot{a} + 1 &= -\kappa\left(p - \frac{1}{4}\kappa s^2\right)a^2 \end{aligned}$$

$$s^2 = \frac{1}{8}(\hbar cn)^2$$

Spin and torsion modify the energy density with a **negative** term proportional to the square of the fermion number density, which acts like **repulsive gravity** and prevents the scale factor from reaching zero. The Big Bang singularity is avoided.

# Universe with spin fluid

For relativistic matter, Friedman equations can be written in terms of temperature.

$$\frac{\dot{a}^2}{c^2} + k = \frac{1}{3}\kappa\tilde{\epsilon}a^2 = \frac{1}{3}\kappa(h_*T^4 - \alpha h_{nf}^2T^6)a^2,$$

$$\alpha = \kappa(\hbar c)^2/32$$

$$\frac{\dot{a}}{a} + \frac{\dot{T}}{T} = \frac{cK}{3h_{n1}T^3},$$

$$K = \beta(\kappa\tilde{\epsilon})^2,$$

2<sup>nd</sup> Friedman equation is rewritten as 1<sup>st</sup> law of thermodynamics for constant entropy. Parker-Starobinskii-Zel'dovich particle production rate  $K$ , proportional to the square of curvature, produces entropy in the Universe. No reheating needed.

# Generating inflation with only 1 parameter

Near a bounce:

$$\frac{\dot{a}}{a} \left[ 1 - \frac{3\beta}{c^3 h_{n1} T^3} \left( \frac{\dot{a}}{a} \right)^3 \right] = -\frac{\dot{T}}{T}$$

To avoid eternal inflation:

$$\beta < \beta_{\text{cr}} = \frac{\sqrt{6} h_{n1} h_{nf}^3 (\hbar c)^3}{32 h_*^3} \approx \frac{1}{929}$$

During an expansion phase, near critical value of particle production coefficient  $\beta$ :

$$\dot{T} \approx 0$$

$$\frac{\dot{a}}{a} \approx \frac{c\beta(\kappa\tilde{\epsilon})^2}{3h_{n1}T^3} \approx c \left( \frac{1}{3} \kappa\tilde{\epsilon} \right)^{1/2}$$


$$\tilde{\epsilon} \approx \frac{h_*^3}{8\alpha^2 h_{nf}^4}$$

**Exponential expansion** lasts about  $\tau = \frac{\alpha h_{nf}^2}{c} \left( \frac{3}{\kappa h_*^3} \right)^{1/2}$  then  $T$  decreases.

Torsion becomes weak and radiation dominated era begins.  
No scalar fields needed.



If quantum effects in the gravitational field near a bounce produce enough matter, then the closed Universe can reach a size at which dark energy becomes dominant and expands to infinity.




Otherwise, the Universe contracts to another bounce (with larger scale factor) at which it produces more matter, and expands again.

The temperature at a bounce depends on the number of elementary particles and the Planck temperature.

Numerical integration of the equations shows that the numbers of bounces and e-folds depend on the particle production coefficient but are not too sensitive to the initial scale factor.




$\beta/\beta_{cr}$	Number of bounces
0.996	1
0.984	2
0.965	3
0.914	5
0.757	10

The Universe might have originated from the interior of a black hole. Accordingly, every black hole may create a new universe on the other side of its event horizon and become a wormhole to that universe (Novikov, Pathria, Smolin, NP, and others).





# Dynamics of the early Universe

$$\beta/\beta_{\text{cr}} = 0.9998$$



It is possible to find a scalar field potential which generates a given time dependence of the scale factor, and calculate the parameters which are being measured in CMB.



Consistent with Planck 2015.

S. Desai & NP, PLB 755, 183 (2016)

## Acknowledgments:

University of New Haven & University Research Scholar program at UNH

Dr. Shantanu Desai

## Torsion research at CfA:

George Field, Rosanne di Stefano

## References:

[arXiv.org > gr-qc > arXiv:1410.3881](https://arxiv.org/abs/1410.3881)

Kibble, JMP 2, 212 (1961)

Sciama, RMP 36, 463 (1964)

Hehl, von der Heyde & Kerlick, PRD 10, 1066 (1974)

Ellis & Madsen, CQG 8, 667 (1991)

NP, PLB 694, 181 (2010); PRD 85, 107502 (2012)

Desai & NP, PLB 755, 183 (2016)